arXiv:1205.1821v1 [physics.space-ph] 8 May 2012

Non-gaussian statistics, maxwellian derivation and stellar polytropes

E. P. Bento[1],[*] J. R. P. Da Silva[2],[†] and R. Silva[1,2‡]

[1] *Universidade Federal do Rio Grande do Norte,*
UFRN, Departamento de Física C. P. 1641,
Natal – RN, CEP 59072–970, Brazil and
[2] *Universidade do Estado do Rio Grande do Norte, UERN,*
Departamento de Física, Mossoró – RN, CEP 59610–210 , Brazil

(Dated: March 3, 2013)

Abstract

In this letter we discuss the Non-gaussian statistics considering two aspects. In the first, we show that the Maxwell's first derivation of the stationary distribution function for a dilute gas can be extended in the context of Kaniadakis statistics. The second one, by investigating the stellar system, we study the Kaniadakis analytical relation between the entropic parameter κ and stellar polytrope index n. We compare also the Kaniadakis relation $n = n(\kappa)$ with $n = n(q)$ proposed in the Tsallis framework.

PACS numbers: 05.45.+b; 05.20.-y; 05.90.+m

---

[*] Electronic address: eliangela@dfte.ufrn.br
[†] Electronic address: joseronaldo@uern.br
[‡] Electronic address: raimundosilva@dfte.ufrn.br

I. INTRODUCTION

The nonextensive statistical mechanics (NSM) [1] and extensive generalized power-law statistics [2] are based on the mathematical generalization of the Boltzmann exponential distribution, namely:

$$f \sim \exp(-Energy/Thermal \quad energy). \tag{1}$$

Whereas in [1], the power-law distribution, $f \sim [1-(1-q)Energy/Thermal \quad energy]^{1/(1-q)}$, obeys the Tsallis distribution of NSM [1], the so-called κ-statistics [2] provides a power-law distribution function and the κ-entropy which emerges in the context of the special relativity and in the *kinetic interaction principle*. Formally, the κ-framework is defined by considering the expressions

$$\exp_\kappa(f) = (\sqrt{1+\kappa^2 f^2} + \kappa f)^{1/\kappa}, \tag{2}$$

$$\ln_\kappa(f) = \frac{f^\kappa - f^{-\kappa}}{2\kappa}, \tag{3}$$

where the κ-entropy associated with the κ-statistics is given by [2]

$$S_\kappa = -\int d^3p f \ln_\kappa f = -\langle \ln_\kappa(f) \rangle. \tag{4}$$

The expressions above reduces to the standard results in the limit $\kappa = 0$.

The κ-statistics leads to a class of one parameter deformed structures with interesting mathematical properties [3], as well as a connection with the generalized Smoluchowski equation [4] and the relativistic nuclear equation of state for nuclear matter [5]. A fundamental test, the so-called Lesche stability was also studied in the context of κ-framework [6]. It was also shown that it is possible to obtain a consistent form for the entropy which is linked with a two-parameter deformations of logarithm function and generalizes the Tsallis, Abe and Kaniadakis logarithm behaviours [7]. There exist some applications related with the κ-statistic, namely, cosmic rays flux, rain events in meteorology [3], quark-gluon plasma [8], kinetic models describing a gas of interacting atoms and photons [9], fracture propagation phenomena [10], and construct financial models [11]. In the theoretical front, some studies on the canonical quantization of a classical system has also been investigated [12], as well as H-theorem in relativistic and non-relativistic domain [13]

In the astrophysical domain, the first application has been the simulation in relativistic plasmas. In this regard, the power-law energy distribution provides a strong argument in favour of the Kaniadakis statistics [14]. Additionally, the viability of the non-gaussian statistics has been investigated in stellar astrophysics viewpoint: the distributions of projected rotational velocity measurements of stars in the Pleiades open cluster [15] and in the main sequence field stars [16], as well as the strong dependence between the stellar-cluster ages and the power-law distributions [17].

The aim of this letter is twofold. First, to show that the Maxwell's first derivation of the stationary distribution function for a dilute gas can be extended in the context of Kaniadakis statistics; Second, considering the principle of maximum entropy for a stellar self-gravitating system, we investigate an analytical relation between entropic parameter κ and stellar polytrope index n. It is also show that the function $n = n(\kappa)$ has a similar behaviour to Tsallis expression $n = n(q)$ [18].

This letter is organized as follows. In Section 2, we show the correspondence between the κ-statistics introduced by Kaniadakis and the velocity distribution for a Maxwellian gas, by assuming a Non-gaussian generalization of the separability hyphothesis originally proposed by Maxwell [19]. In Section 3, we discuss a connection between the Kaniadakis statistics and polytropic index in the context of the self-gravitating system and we compare our results with ones studied in Ref. [18]. Finally, Section 4 is devoted to conclusions and discussion.

II. NON-GAUSSIAN MAXWELLIAN DISTRIBUTION FUNCTION

In this section, in order to introduce the generalization of the Maxwell distribution in the context of Kaniadakis framework, let us consider a spatially homogeneous gas, supposed in equilibrium (or in stationary state) at temperature T, in such a way that $F(v)d^3v$ is the number of particles with velocity in the volume element d^3v around v. In Maxwell's derivation, the 3-dimensional distribution is factorized (lottery assumption) and depends only on the magnitude of the velocity [19, 20]

$$F(v)d^3v = f(v_x)f(v_y)f(v_z)dv_x dv_y dv_z \quad , \tag{5}$$

where $v = \sqrt{v_x^2 + v_y^2 + v_z^2}$ and $F(v)$ is the standard Maxwellian distribution function, given by

$$F(v) = A\exp(-mv^2/k_BT), \tag{6}$$

where $A = \left(\frac{m}{2\pi k_BT}\right)^{3/2}$ is the normalization constant.

As a matter of fact, in the κ-statistic context described by (4), the starting basic hypothesis (5), which takes into account the isotropy of all velocity directions, must somewhat be modified. In statistical viewpoint, Maxwell's ansatz assume that the three components of the velocity are uncorrelated. However, this property does not hold in the systems endowed with long range interactions, or statistically correlated, where Kaniadakis distribution has been observed [15, 16]. Notice that the Maxwell ansatz is equivalent to express $\ln F$ as the sum of the logarithms of the one dimensional distribution functions associated with each velocity component. A simple and natural way to generalize this procedure within the Kaniadakis framework, in order to introduce correlations between the velocity components, would be to replace the logarithm function by a power law. However, in order to recover the ordinary logarithmic ansatz as a particular limiting case, it is convenient to express the power generalization in terms of the function $\ln_\kappa$ defined by Eq. (3), which is a combination of a power function plus appropriate constants. Mathematically, the consistent κ-generalization of (5) is given by

$$\begin{aligned} F\left(\sqrt{v_x^2+v_y^2+v_z^2}\right) d^3v \; &= \; \exp_\kappa[\ln_\kappa f(v_x) + \ln_\kappa f(v_y) + \\ & \quad \ln_\kappa f(v_z)] dv_x dv_y dv_z \;, \end{aligned} \tag{7}$$

where the κ-exponencial and κ-logarithm are given by identities (2) and (3). In particular, in the limit $\kappa = 0$ the standard expression (5) is recovered. Note also that $\ln_\kappa[\exp_\kappa(f)] = \exp_\kappa[\ln_\kappa(f)] = f$, and $\frac{d\ln_\kappa(f)}{dx} = \frac{f^\kappa + f^{-\kappa}}{2f}\frac{df}{dx}$ are satisfied. The logarithmic derivative of (7) with respect to v_i is

$$\begin{aligned} \frac{\partial}{\partial v_i} \ln_\kappa F\left(\sqrt{v_x^2+v_y^2+v_z^2}\right) \; &= \; \frac{\partial}{\partial v_i} \ln_\kappa\{\exp_\kappa[\ln_\kappa f(v_x) + \\ & \quad \ln_\kappa f(v_y) + \ln_\kappa f(v_z)]\}, \end{aligned} \tag{8}$$

with $i = x,y,z$.

Using the properties above mentioned we can write

$$\frac{F^\kappa + F^{-\kappa}}{2F}\frac{\partial}{\partial v_i} F\left(\sqrt{v_x^2+v_y^2+v_z^2}\right) = \frac{\partial}{\partial v_i}[\ln_\kappa f(v_i)] \;\; . \tag{9}$$

Equivalently,

$$\frac{F^\kappa + F^{-\kappa}}{2F}\frac{F'(\chi)}{\chi} = \frac{1}{v_i}\frac{\partial}{\partial v_i} \ln_\kappa f(v_i) \;\; , \tag{10}$$

where $\chi = \sqrt{v_x^2+v_y^2+v_z^2}$. Now, defining

$$\Phi(\chi) = \frac{F^\kappa + F^{-\kappa}}{2F}\frac{F'(\chi)}{\chi} \;\; , \tag{11}$$

the equation(10) can be written as

$$\Phi(\chi) = \frac{1}{v_i}\frac{d}{dv_i}[\ln_\kappa f(v_i)] \ . \tag{12}$$

The second member of Eq. (12) has dependence only on v_i, with $i = x,\ y,\ z$. Hence, the Eq. (11) can be satisfied only if all its members are equal to one and the same constant, not depending on any of the velocity components. Thus we can make $\Phi(\chi) = -m\gamma$, obtaining

$$\frac{1}{v_i}\frac{d}{dv_i}[\ln_\kappa f(v_i)] = -m\gamma \ , \tag{13}$$

where $\gamma = \frac{1}{k_B T}$.

In general, we have

$$\ln_\kappa f(v_i) = -\frac{m\gamma v_i^2}{2} \ , \tag{14}$$

which provides

$$f(v_i) = \frac{1}{Z}\exp_\kappa\left(-\frac{mv_i^2}{2k_B T}\right) \ , \tag{15}$$

where $1/Z$ is the κ-normalization constant.

In order to calculate the so-called κ-normalization, let us introduce the expression [2]

$$Z = \int_{\mathcal{R}} d^n v \exp_\kappa\left(-\frac{mv^2}{2k_B T}\right) \ , \tag{16}$$

where n (=1,2,3) is the number from degree of freedom. Here, considering $n = 1$, $a = \frac{m}{2k_B T}$ and $x = av^2$, we obtain

$$Z = \frac{1}{\sqrt{a}}\int_0^\infty x^{-1/2}\exp_\kappa(-x)dx. \tag{17}$$

Using the generalized gamma functions [2]

$$\begin{aligned} \int_0^\infty x^{r-1}\exp_\kappa(-x)dx \ &= \ \frac{[1+(r-2|\kappa|)|2\kappa|^r]}{[1-(r-1)|\kappa|]^2-\kappa^2} \\ &\quad \times \frac{\Gamma\left(\frac{1}{|2\kappa|}-\frac{r}{2}\right)}{\Gamma\left(\frac{1}{|2\kappa|}+\frac{r}{2}\right)}\Gamma\left(\frac{1}{2}\right) , \end{aligned} \tag{18}$$

and after some algebra, we obtain

$$Z = \sqrt{\frac{\pi k_B T}{m}}\frac{|\kappa|^{-1/2}}{\left[1-\frac{1}{2}|\kappa|\right]}\frac{\Gamma\left(\frac{1}{|2\kappa|}-\frac{1}{4}\right)}{\Gamma\left(\frac{1}{|2\kappa|}+\frac{1}{4}\right)} \ . \tag{19}$$

It is easy to see that the standard Maxwellian result $(1/Z) = (m/2\pi k_B T)^{1/2}$ is recovered for gaussian limit $\kappa = 0$.

In order to calculate the complete distribution, we insert the expressions (15) into of Eq. (7) for obtain

$$F(v) = \frac{1}{Z}\left[\sqrt{1+\kappa^2\left(-\frac{mv^2}{2k_BT}\right)^2} - \frac{\kappa m v^2}{2k_BT}\right]^{\frac{1}{\kappa}} . \tag{20}$$

Using the expression (16) for $n = 3$, we can show that the normalization for the complete distribution is given by

$$\frac{1}{Z} = \left(\frac{m|\kappa|}{\pi k_B T}\right)^{3/2}\left[1+\frac{3}{2}|\kappa|\right]\frac{\Gamma\left(\frac{1}{|2\kappa|}+\frac{3}{4}\right)}{\Gamma\left(\frac{1}{|2\kappa|}-\frac{3}{4}\right)} , \tag{21}$$

As expected, the κ-distribution above is isotropic meaning that all velocity directions are also equivalent in this generalized context. Here, we emphasize that in the limit $\kappa = 0$ both the normalization and the distribution function reproduces the standard Maxwellian (6).

III. NON-GAUSSIAN FRAMEWORK AND STELLAR POLYTROPES

As mentioned in the introduction, we shall discuss a connection between the Kaniadakis statistics and polytropic index in the context of the self-gravitanting system. Following a procedure considered in Ref. [18], let us start with the Kaniadakis generalized entropy of index κ of the form

$$S_\kappa(f) = -\int \frac{f^{1+\kappa} - f^{1-\kappa}}{2\kappa} d\Omega, \tag{22}$$

where $d\Omega = d^3\boldsymbol{r}d^3\boldsymbol{v}$, and the parameter $\kappa \neq 0$ provides a possible generalization of the Boltzmann Gibbs entropy. The extremum entropy state can be derived by varying S_κ with respect to f. Using the Lagrange multipliers α and β, the extremum solution subject to constraints $M = \int f d\Omega$ and $E = K + U = \frac{1}{2}\int v^2 f d\Omega + \frac{1}{2}\int \phi d\Omega$ [1] is obtained from

$$\delta S_\kappa - \alpha\delta M - \beta\delta E = 0 \tag{23}$$

---

[1] The mass and energy total of the self gravitating system governed by the Vlasov and Poisson equations. For more details, see Ref. [18]

which leads to

$$\int\left\{-\frac{1}{2\kappa}\left[(1\,+\,\kappa)f^{\kappa}-(1-\kappa)f^{-\kappa}\right]-\alpha\right.$$
$$\left.-\beta\left(\frac{v^2}{2}+\phi\right)\right\}\delta f d\Omega=0. \tag{24}$$

We have used the relation $\int\delta\phi f d\Omega=\int\phi\delta f d\Omega$ for derivation of above expression. Here, ϕ denotes gravitational potential, and the equation (23) must be satisfied independently of the choice of δf. Thus, we obtain the following distribution function

$$f(\boldsymbol{r},\boldsymbol{v})=B\left[\phi_0(\boldsymbol{\epsilon})-\phi(\boldsymbol{r})-\frac{v^2}{2}\right]^{1/\kappa} \tag{25}$$

where the constants B and ϕ_0 are defined by

$$B=\left(\frac{\kappa\beta}{(1+\kappa)}\right)^{1/\kappa}\quad,$$

$$\phi_0(\epsilon)=\frac{-\sqrt{\kappa^2(-\alpha-\beta\epsilon)^2+1+\kappa^2}-\alpha\kappa}{\beta\kappa}. \tag{26}$$

Now, comparing the distribution (25) with the polytropic spheres distribution $f\sim\varepsilon^{n-3/2}$ (ϵ is the relative energy of a star) [21], we obtain the relation between the stellar polytropic index and the entropic index, given by

$$n=\frac{3}{2}+\frac{1}{\kappa}. \tag{27}$$

In order to compare the expression $n=n(\kappa)$ with the Tsallis relation $n=n(q)$, we present the expression discussed in Ref. [18]

$$n=\frac{3}{2}+\frac{1}{q-1}, \tag{28}$$

which include the isothermal situation for $n\to\infty$, i.e., the Gaussian limit $q=1$. In order to guarantee the conservation of mass and energy, the Tsallis' parameter q should be larger than $9/7$.

In Figure 1 we display the dependence between the entropic indexes $(\kappa;q)$ and the polytropic index (n), given by Eqs. (27) and (28), respectively. We show that the polytropic index diverge for the non-gaussian parameters $\kappa=0, q=1$, i.e., the isothermal spheres separates the polytropes into two branches. For $\kappa>0$ and $q>1$, n ranges from ∞ to $3/2$. From $f\sim\varepsilon^{n-3/2}$, we see that ε is decreasing function of the energy and, when $n=3/2$,



FIG. 1: Polytrope index n as a function of the Kaniadakis (κ) and Tsallis (q) parameters. The vertical dotted lines mark the limits of validity of κ-statistical ($-1 \leq \kappa \leq 1$) [2].

the distribution function becomes a constant independent of energy. On the other hand, it is well known that for any astrophysical system, n should be positive and higher than 1/2 [21]. Thus the intervals $0 \leq q \leq 1$ and $-1 \leq \kappa \leq 0$ represent forbidden regions, since the values of n index tend to be smaller than 1/2 and negative.

It is also worth observing that for $\kappa < -1$ and $q < 0$ also provides $n > 1/2$, however this limits on $\kappa; q$ violates the validity of κ-statistics [2] and the non-extensive H-theorem [23], respectively. As is well known, for the polytropic index $n > 5$, the density falls off so slowly at large radii that the mass is infinite [21]. Figure 1 shows that only the branches of the right-side are physically significant for $n < 5$. Therefore, the physical values of the Tsallis parameter q should be larger than 9/7[2], and the Kaniadakis parameter κ should be constrained to interval of validity $\kappa \in [2/7; 1]$ represented by the lower dashed rectangle.

---

[2] Investigating a collisionless stellar gas in the context from Non-extensive Kinetic Theory, the bound $q > 9/7$ also was calculated in Ref. [22]

IV. CONCLUSIONS

In this work we have studied the Kaniadakis statistic based on the generalized Maxwellian formulation for the κ-statistic and, as one application, we investigate the physical effect on the stellar polytropic system. In the first part, we conclude that there is a Kanidakis velocity distribution given by Eq. (20), that is uniquely determined by the requirements of (i) isotropy and (ii) a suitable generalization of the factorizability condition. From the physical viewpoint, this generalization introduces statistical correlations between the velocity components, when we replace the logarithm function by a power law (similar argument was considered in Refs. [23, 24] for the non-gaussian generalization of molecular chaos hypothesis). In particular, Maxwell expressions are recovered in the gaussian limit, $\kappa \to 0$.

From an application perspective, we have shown that the expressions for Tsallis and Kaniadakis, given by Eqs. (28) and (27), present similar behaviours. However, the astrophysical limit on the polytropic index $1/2 < n < 5$, provides the constrains $\kappa \in [2/7; 1]$ and $q > 9/7$ for the Kaniadakis and Tsallis parameters, respectively. It is worth mentioning that the gaussian limit $\kappa = 0$, equivalent the Tsallis expression Eq. (28) for $q = 1$, reproduces Maxwellian isothermal spheres or, equivalently, $n = \infty$.

Acknowledgments

E.P.B. and RS acknowledges financial support from CNPq.

---

[1] C. Tsallis, J. Stat. Phys. **52**, 479 (1988)

[2] G. Kaniadakis, Physica A **296**, 405 (2001); Phys. Rev. E **66**, 056125 (2002); Phys. Rev. E **72**, 036108 (2005); Eur. Phys. J. B **70** (2009)(special number); Europhys. Lett. **92**, 35002 (2010); Phys. Lett. A **375**, 356 (2011)

[3] G. Kaniadakis, Phys. Lett. A **288**, 283 (2001); G. Kaniadakis, A.M. Scarfone, Physica A **305**, 69 (2002); G. Kaniadakis, P. Quariti, A.M. Scarfone, Physica A **305**, 76 (2002)

[4] P. H. Chavanis, P. Laurencot, M. Lemou, Physica A **341**, 145 (2004)

[5] F. I. M. Pereira, R. Silva, and J. S. Alcaniz, Nuc. Phys. A **828**, 136 (2009)

[6] G. Kaniadakis, A.M. Scarfone, Physica A **340**, 102 (2004); G. Kaniadakis, M. Lissia, A.M. Scarfone, Physica A **340**, 41 (2004)

[7] G. Kaniadakis, M. Lissia, A. M. Scarfone, Phys. Rev. E **71**, 046128 (2005)

[8] A. M. Teweldeberhan, H. G. Miller, R. Tegen, Int. J. Mod. Phys. E **12**, 669 (2003)

[9] A. Rossani, A. M. Scarfone, J. Phys. A: Math. Gen. **37** 4955 (2004)

[10] M. Cravero, G. Labichino, G. Kaniadakis, E. Miraldi, A. M. Scarfone, Physica A **340**, 410 (2004)

[11] D. Rajaonarison, D. Bolduc, H. Jayet, Econ. Lett. **86** 13 (2005)

[12] A. M. Scarfone, Phys. Rev. E **71**, 051103 (2005)

[13] R. Silva, Eur. Phys. J. B **54**, 499 (2006); Phys. Lett. A **352** 17 (2006)

[14] G. Lapenta, S. Markidis, A. Marocchino and G. Kaniadakis, Astrophys. Journ. **666**, 949 (2007)

[15] J. C. Carvalho, R. Silva, J. D. Jr. do Nascimento, J. R. De Medeiros, Europhys. Lett. **84**, 59001 (2008)

[16] J. C. Carvalho, J. D. Jr. do Nascimento, R. Silva, J. R. De Medeiros, Astrophys. Journ. Lett. **696**, L48 (2009)

[17] J. C. Carvalho, R. Silva, J. D. Jr. do Nascimento, B. B. Soares and J. R. De Medeiros, Europhys. Lett. **91**, 69002 (2010)

[18] A. Plastino, A. R. Plastino, Phys. Lett. A **174**, 384 (1993); A. R. Plastino, Europhysics News **36**, 208 (2005)

[19] J. C. Maxwell, Philos. Mag. Ser. 4, **20**, 21 (1860)

[20] A. Sommerfeld, Thermodynamics and Statistical Mechanics, Lectures on Theorethical Physics, Vol. V, Academic Press, New York (1993)

[21] J. Binney and S. Tremaine, Galactic Dynamics, Princeton University Press, Princeton (1987)

[22] J. A. S. Lima, R. E. de Souza, Physica A **350**, 303 (2005)

[23] J. A. S. Lima, R. Silva, A. R. Plastino, Phys. Rev. Lett. **86**, 2938 (2001); R. Silva, J. A. S. Lima, Phys. Rev. E **72**, 057101 (2005); R. Silva, D. H. A. L. Anselmo, J. S. Alcaniz, Europhys. Lett. **89**, 10004 (2010); R. Silva, D. H. A. L. Anselmo, J. S. Alcaniz, Europhys. Lett. **89**; 59902 (2010) (Erratum)

[24] A. P. Santos, R. Silva, J. S. Alcaniz, D. H. A. L. Anselmo, Phys. Lett. A **375**, 352 (2011)